

## Bibliography

1. American College of Obstetricians and Gynecologists (2009). Management of stillbirth. ACOG Practice Bulletin, 102.
2. Stacey, T., Thompson, J., Mitchell, E.A., Ekeroma, A.J., Zuccollo, J.M., & McCowan, L.M.E. (2011). Relationships between obesity, ethnicity and risk of late stillbirth: a case control study. BMC Pregnancy and Childbirth, 11 (3). Doi 10.1186/1471-2393-11-3
3. Pasupathy, D., & Smith, G.C.S. (2005). The analysis of factors predicting antepartum stillbirth. Minerva Ginecologica: 57 (4), p 397-410.
4. McGregor, J. (2011). Prenatal-onset Group B Strep Disease: Giving a name to Group B Strep Disease. Obtained online: [www.obgyn.net/pregnancy-birth/pregnancy-birth.asp](http://www.obgyn.net/pregnancy-birth/pregnancy-birth.asp).
5. Wilson, B. (2008). Sonography of the placenta and umbilical cord. Radiologic Technology, 79(4): 333S-345S.
6. Azpurua, H., Funai, E.F., Coraluzzi, L.M., Doherty, L.F., Sasson, I.E., Kliman, M., & Kliman, H.J. (2010). Determination of placental weight using two-dimensional ultrasonography and volumetric mathematic modeling. American Journal of Perinatology, 27(2): 151-154.
7. Royal College of Obstetricians and Gynaecologists (2011). Green-top Guideline 57: Reduced Fetal Movements.
8. Stillbirth Collaborative Research Network Writing Group (2011). Causes of death among stillbirths. JAMA, 306 (22): 2459-2468.
9. Warland, J., McCutcheon, H., & Baghurst, P. (2008). Maternal blood pressure in pregnancy and stillbirth: A case-control study of third-trimester stillbirth. Am J Perinatol; 25: 311-317.
10. Stacey, T., Thompson, J.M.D., Mitchell, E.A., Ekeroma, A.J., Zuccollo, J.M., & McCowan, L.M.E. (2011). Association between maternal sleep practices and risk of late stillbirth: a case-control study. BMJ: doi: 10.1136/bmj.d3403.
11. Hasegawa, J., Matsuoka, R., Ichizuka, K., Sekizawa, A., & Okai, T. (2008). Ultrasound diagnosis and management of umbilical cord abnormalities. ObstetGynecol; 48(1); 23-27.
12. International Society for the Study and Prevention of Perinatal and Infant Death (2012). The Role of Fetal Movement Counting in Reducing Stillbirth. Online: [http://www.ispid.org/fetalmovement2.html?&no\\_cache=1&word\\_list%5B0%5D=kick&word\\_list%5B1%5D=counting&type=98](http://www.ispid.org/fetalmovement2.html?&no_cache=1&word_list%5B0%5D=kick&word_list%5B1%5D=counting&type=98).
13. Edwards N, Sullivan CE. Sleep-disordered breathing in pregnancy. Sleep Medicine Clinics 2008;3:81-95
14. Lee KA, Gay CL. Sleep in late pregnancy predicts length of labor and type of delivery. Am J Obstet Gynecol. 2004 Dec;191(6):2041Qiu C,
15. Enquobahrie D, Frederick IO, Abetew D, Williams MA. Glucose intolerance and gestational diabetes risk in relation to sleep duration and snoring during pregnancy: a pilot study. BMC Womens Health. 2010 May 14;10:17

To learn more about See Me Feel Me, please visit [www.starlegacyfoundation.org](http://www.starlegacyfoundation.org)

This information is not intended to replace medical advice. Please contact your health care provider with any questions or concerns.

See Me, Feel Me is an education project of

The Star Legacy Foundation.





Pregnancy is an exciting and beautiful time when parents prepare to welcome a new life into their family. The development of this life is nothing short of a miracle; but, unfortunately (like most things in life) there are no guarantees and sometimes things can go wrong. Nearly 30,000 babies are stillborn\* in the United States every year.<sup>1</sup> The good news is there are measures you can take to enhance your chances of having a healthy baby.

### **Your parenting duties begin now!**

#### **WHY SEE ME, FEEL ME?**

First – it’s fun! Getting to know your baby is one of the most magical parts of being pregnant! **ONLY YOU** get to have this amazing experience to know your child before he/she is born! *This is a great privilege, but also a great responsibility.*



Secondly, some conditions mentioned in this brochure may increase the risk of adverse pregnancy outcomes including premature labor and delivery, birth injury, and stillbirth. <sup>(7, 8)</sup> Having as much information about you, your baby, and your baby’s environment as possible allows you and your healthcare team to make the best decisions to improve chances of your baby’s healthy arrival. Finally, we believe the mother knows her baby best!

Mom’s intuition and concerns about unusual symptoms should play a significant role in the care of any pregnancy.

\*stillbirth is defined as the death of an unborn baby at 20 or more weeks gestation.

#### **OTHER CONSIDERATIONS**

New research is being conducted every day to learn more about reducing the risk of poor pregnancy outcomes. The following topics are emerging as possible benefits. Discuss these with your obstetric care provider.

- ◇ High blood pressure is known to cause problems, but a recent study indicates low blood pressure may be of concern as well. If you have low or borderline low blood pressure, discuss this risk with your healthcare team. <sup>(9)</sup>
- ◇ If you snore or have trouble breathing at night, discuss this with your healthcare provider as sleep apnea has been linked to high blood pressure and pre-eclampsia, and it can be treated, even during pregnancy.<sup>(13)</sup>
- ◇ Do your best to maintain a healthy sleep schedule. Poor sleep is associated with a greater risk of longer labors and c-sections. Tell your provider if sleep is a concern. <sup>(14)</sup>
- ◇ Sleeping on your left side may help reduce the risk of stillbirth. Doctors, nurses, and midwives often use lying on the left side to enhance blood flow to the baby during labor. A new study shows this may also be true throughout the pregnancy. <sup>(10)</sup>
- ◇ Invasive exams such as internal (vaginal) exams and membrane stripping are being shown to increase the risk of infection spreading to the uterus and baby. Discuss the risks and benefits of any invasive procedure with your provider before it is conducted. <sup>(4)</sup>
- ◇ Extremes in maternal age (less than 20 years old or older than 35 years old) may increase the risk of poor outcomes. Also, African-American women experience higher rates of stillbirth than women of other ethnicities in the United States. African-American women, women less than 20 years of age, and women over 35 years of age should discuss this risk with their providers to determine if additional testing or surveillance is needed. <sup>(8)</sup>
- ◇ Women having their first baby or those who have had a previous stillbirth, premature birth, or birth injury have a higher risk of stillbirth. If this is your first pregnancy or you have experienced these issues with a previous pregnancy, ask your provider about this increased risk and what to watch for. <sup>(8)</sup>

## WHAT HAPPENS NEXT



When you report any changes or concerns to your health care provider, you should be evaluated further. This may include a non-stress test for 30-60 minutes or more, ultrasound, biophysical profile, daily monitoring, additional visits, or other testing and interventions as determined to be appropriate by your health care provider. Other factors will also be taken into consideration such as your age, blood pressure problems, your family history, and what has happened with your previous pregnancies. (7, 9, 10, 11)

If your prenatal testing and kick counting show a healthy mom, baby, cord, and placenta, that is wonderful! Continue with the kick counting and follow the advice of your health care team. Be aware of how you are feeling and continue to report any changes right away. Ask your healthcare providers to keep you up-to-date about local outbreaks of infection and how to prevent many types of infections. If at any time you experience abdominal or back pain, fever, vaginal bleeding or leaking, contact your health care team immediately. (7)

You are your baby's voice. Be his/her advocate! If you ever don't feel right about your pregnancy or the response to your concerns, don't hesitate to ask more questions or obtain a second opinion if necessary.

## BEFORE PREGNANCY

### **1. Visit your healthcare provider before you get pregnant.**

The first step to having a healthy baby is to start with a healthy mother. Schedule an appointment with your healthcare provider to determine if you have any health concerns that should be addressed before or during pregnancy. Be sure to mention your intentions to become pregnant. Your provider may evaluate your immune status, screen for common infections, or test for other health indicators before you become pregnant. Certain vitamins or supplements, like folic acid, may be recommended to ensure you and your baby's health. Many common medications (whether prescribed, natural, or over the counter) and diseases like diabetes, autoimmune disorders, heart disease, thyroid disorders, alterations in sleep, hypertension, infections, renal disease, and others can predispose you to complications during the pregnancy.<sup>(1)</sup> The risk of miscarriage, stillbirth, premature birth, and birth injury can be reduced with good attention to the health of you and your baby. Your health care provider will help you develop a plan that is best for you at this time.



### **2. If you smoke or use recreational drugs, it is time to quit. Reduction/elimination of alcohol is recommended.**

Smoking, alcohol use, and drug use have been proven to be harmful to you and your baby. <sup>(1)</sup>Your health care provider may be able to recommend specific programs to assist your efforts to quit if needed.

### **3. Adopt a healthy lifestyle.**

If you are overweight, now is the time to begin a weight-loss program to include a healthy diet and active lifestyle. Overweight and obese women have a higher risk of complications during pregnancy including high blood pressure, diabetes, and stillbirth.<sup>(2)</sup> Weight loss during pregnancy generally is not recommended. Discuss a healthy weight and nutrition plan with your health care provider. If you are often stressed, talk to your provider about reducing or eliminating stress in your life before and during pregnancy.

